

West Lothian Health and Wellbeing Profiles – key indicators and overview

Cite as: Millard A, McCartney G, MacKinnon A, Van Heelsum A, Gasiorowski A, Barkat S. *West Lothian Health and Wellbeing Profiles – key indicators and overview*. Edinburgh: ScotPHO; 2016.

Introduction

The ScotPHO Health and Wellbeing profiles (www.scotpho.org.uk/comparative-health/profiles/online-profiles-tool) provide a set of health and wellbeing indicators for Scottish local authority areas. The indicators are displayed in spine charts for each local authority and are measured in the same way in each part of Scotland. One of the key aims of the profiles is to allow comparisons between areas, and the data that are available demonstrate the inequalities in outcomes between areas.


Feedback shows that some people find using the online profiles tool daunting, therefore this report will provide key points without the need to go into the tool. For further details or up-to-date figures see the final section of this report, which explains how to use the tool.

The featured spine chart compares the local and national position for each indicator. Also included – for the worst key indicator locally – are a trend chart, showing changes over time in comparison with Scotland, and a rank chart, showing how your local authority compares with the other 31 local authorities. These charts can be accessed online by clicking on an individual indicator name in the spine chart.

Throughout this report we refer to the European Age Standardised Rate (EASR), which is the rate per 100,000 population adjusted for the different age balances. Unless the indicator is given for one sex only, the EASRs adjust for both differing age and sex balance. Crude rates are the rates before any adjustment. Unless specified as crude, 'rate' always means the EASR in this report.

Key differences from Scotland in health and wellbeing indicators for West Lothian

The 'worst' indicators, ranked locally are reported below in the key differences summary. The aim is to provide an indication of which areas might be considered as priorities for improvement.



Methods

Indicators are highlighted only when it is considered that the difference between the local and national figure is not just due to random variation (in other words the difference is statistically significant). Figures were ordered from the worst to the best, then the six worst were selected and compared to the Scottish averages. This information may be of use in health improvement planning, but must be considered in a local context.

The information in this report is a snapshot taken on a particular date (25 February 2016). The indicators are updated continuously, and therefore more up-to-date information may be available online. To avoid disclosure as a result of small numbers, many indicators use aggregated figures where necessary, for example over a three- or five-year period, such as 2012–14. The data can be collected for a calendar year, such as 2012, or a financial year, such as 2012/13.

Key indicators for West Lothian

For West Lothian in 2012/2013, the percentage of active travel to work was 10%, which was 38% lower than the Scottish level of 16%.


In 2012/13–2014/15, the percentage of mothers smoking during pregnancy was 22%, which was 17% higher than the Scottish level of 19%.

In 2011–2013, the rate for patients with a psychiatric hospitalisation was 341/100,000, which was 17% higher than the Scottish level of 292/100,000.

In 2014, the percentage of young people not in employment, education or training (NEET) was 8%, which was 16% higher than the Scottish level of 7%.

In 2011–2013, the rate for patients (65+) with multiple emergency hospitalisations was 5945/100,000, which was 15% higher than the Scottish level of 5159/100,000.

In 2012/13–2014/15, the percentage of babies exclusively breastfed at 6–8 weeks was 24%, which was 12% lower than the Scottish level of 27%.



Health & Wellbeing Profiles (West Lothian)

| Domain | Indicator | Period | Number | Measure | Type | National Average | 'Worst' | Scotland Comparator | 'Best' |
|-----------------------------|--|--------|--------|---------|------|------------------|---------|---------------------|--------|
| Life Expectancy & Mortality | 1 Male life expectancy ¹⁸ | 2011 | n/a | 77.1 | yrs | 76.6 | | | |
| | 2 Female life expectancy ¹⁸ | 2011 | n/a | 80.1 | yrs | 80.8 | | | |
| | 3 Deaths all ages ¹² | 2013 | 1,452 | 1,180.2 | sr4 | 1,169.8 | | | |
| | 4 All-cause mortality among the 15-44 year olds. ¹² | 2013 | 61 | 87.7 | sr4 | 100.5 | | | |
| | 5 Early deaths from CHD (<75) ¹² | 2012 | 89 | 64.4 | sr4 | 60.7 | | | |
| | 6 Early deaths from cancer (<75) ¹² | 2013 | 248 | 157.8 | sr4 | 154.7 | | | |
| Behaviours | 7 Estimated smoking attributable deaths ^{3,13,16} | 2014 | 269 | 345.4 | sr4 | 366.8 | | | |
| | 8 Smoking prevalence (adults 16+) ^{3,14} | 2014 | 41 | 20.6 | % | 20.2 | | | |
| | 9 Alcohol-related hospital stays ¹⁵ | 2014 | 1,112 | 642.2 | sr4 | 671.7 | | | |
| | 10 Deaths from alcohol conditions ¹⁷ | 2012 | 33 | 19.7 | sr4 | 23.1 | | | |
| | 11 Drug-related hospital stays ^{12,15} | 2013 | 226 | 126.6 | sr4 | 122.0 | | | |
| | 12 Active travel to work ^{3,14} | 2013 | 23 | 10.0 | % | 16.0 | | | |
| | 13 Patients registered with cancer ¹² | 2012 | 914 | 646.0 | sr4 | 634.1 | | | |
| | 14 Patients hospitalised with chronic obstructive pulmonary disease (COPD) ^{12,15} | 2012 | 947 | 709.9 | sr4 | 659.9 | | | |
| | 15 Patients hospitalised with coronary heart disease ¹² | 2012 | 599 | 419.7 | sr4 | 440.3 | | | |
| | 16 Patients hospitalised with asthma ¹² | 2012 | 164 | 86.4 | sr4 | 91.2 | | | |
| Ill Health & Injury | 17 Patients with emergency hospitalisations ¹² | 2012 | 13,046 | 8,189.7 | sr4 | 7,500.2 | | | |
| | 18 Patients (65+) with multiple emergency hospitalisations ¹² | 2012 | 1,364 | 5,944.9 | sr4 | 5,159.5 | | | |
| | 19 Road traffic accident casualties ¹² | 2012 | 92 | 53.4 | sr4 | 63.2 | | | |
| | 20 Population prescribed drugs for anxiety/depression/psychosis ³ | 2014 | 31,191 | 17.6 | % | 17.4 | | | |
| | 21 Patients with a psychiatric hospitalisation ¹² | 2012 | 557 | 341.3 | sr4 | 291.6 | | | |
| Mental Health | 22 Deaths from suicide ¹⁷ | 2011 | 25 | 15.0 | sr4 | 14.5 | | | |
| | 23 Adults claiming incapacity benefit/severe disability allowance/employment and support allowance | 2014 | 8,980 | 5.1 | % | 5.1 | | | |
| Social Care & Housing | 24 ³ People aged 65 and over with high levels of care needs who are cared for at home | 2013 | 471 | 38.0 | % | 34.7 | | | |
| | 25 Children looked after by local authority ³ | 2013 | 445 | 10.7 | cr2 | 14.4 | | | |
| Education | 26 Single adult dwellings | 2014 | 27,260 | 35.3 | % | 37.5 | | | |
| | 27 Average tariff score of all pupils on the S4 roll ¹³ | 2012 | n/a | 206.0 | mean | 193.0 | | | |
| | 28 Primary school attendance | 2010 | 13,654 | 95.3 | % | 94.8 | | | |
| | 29 Secondary school attendance | 2010 | 10,464 | 91.9 | % | 91.1 | | | |
| Economy | 30 Working age adults with low or no educational qualifications ³ | 2013 | 12,800 | 11.4 | % | 12.6 | | | |
| | 31 Population income deprived | 2014 | 22,290 | 12.6 | % | 13.1 | | | |
| | 32 Working age population employment deprived | 2014 | 13,490 | 11.7 | % | 12.2 | | | |
| | 33 Working age population claiming Out of Work benefits | 2014 | 13,545 | 11.8 | % | 12.0 | | | |
| | 34 Young people not in employment, education or training (NEET). ³ | 2014 | 650 | 7.5 | % | 6.5 | | | |

| | | | | | | | | |
|--|----|---|------|--------|-------|-----|-------|--|
| | 35 | Children Living in Poverty | 2012 | 6,365 | 14.4 | % | 15.3 | |
| | 36 | People claiming pension credits (aged 60+) | 2014 | 2,550 | 6.9 | % | 6.9 | |
| | 37 | Crime rate | 2014 | 6,237 | 35.2 | cr2 | 40.4 | |
| Crime | 38 | Prisoner population ^{3,13} | 2012 | 171 | 117.2 | sr4 | 171.2 | |
| | 39 | Referrals to Children's Reporter for violence-related offences ³ | 2013 | 51 | 3.0 | cr2 | 2.1 | |
| | 40 | Domestic Abuse ³ | 2014 | 2,218 | 125.2 | cr9 | 112.0 | |
| | 41 | Violent crimes recorded ³ | 2014 | 186 | 10.5 | cr9 | 11.9 | |
| | 42 | Drug crimes recorded ³ | 2014 | 671 | 37.9 | cr9 | 68.9 | |
| | 43 | Population within 500 metres of a derelict site | 2013 | 43,492 | 24.7 | % | 29.7 | |
| Environment | 44 | People living in 15% most 'access deprived' areas | 2014 | 11,900 | 6.7 | % | 15.0 | |
| | 45 | Adults rating neighbourhood as 'a very good place to live' ^{3,14} | 2014 | n/a | 60.8 | % | 55.8 | |
| | 46 | Teenage pregnancies ¹² | 2012 | 220 | 40.9 | cr2 | 41.1 | |
| Women's & Children's Health | 47 | Mothers smoking during pregnancy ¹² | 2013 | 384 | 21.6 | % | 18.5 | |
| | 48 | Low birth weight ¹² | 2013 | 34 | 2.0 | % | 2.0 | |
| | 49 | Babies exclusively breastfed at 6-8 weeks ¹² | 2013 | 466 | 23.6 | % | 26.8 | |
| | 50 | Child dental health in primary 1 | 2013 | 1,383 | 64.3 | % | 66.7 | |
| | 51 | Child dental health in primary 7 | 2013 | 780 | 42.9 | % | 47.7 | |
| | 52 | Child obesity in primary 1 | 2013 | 174 | 8.1 | % | 10.1 | |
| | 53 | Breast screening uptake ¹² | 2011 | 4,969 | 68.0 | % | 72.5 | |
| | 54 | Bowel screening uptake ¹² | 2012 | 13,787 | 52.7 | % | 56.0 | |
| | 55 | Immunisation uptake at 24 months - 5 in 1 ¹² | 2013 | 2,150 | 98.9 | % | 98.2 | |
| | 56 | Immunisation uptake at 24 months - MMR ¹² | 2013 | 2,090 | 96.2 | % | 95.3 | |
| Immunisations and Screening | | | | | | | | |

Key

Notes:

- Data available down to council (local authority) area only.
- Three-year average number, and 3-year average annual measure.
- Indicator based on HB boundaries prior to April 2014.
- Two-year combined number, and 2-year average annual measure.
- All 6 diagnosis codes used in the analysis; please see the technical report for more information.
- Two-year average number, and 2-year average annual measure
- Five-year average number, and 5-year average annual measure
- Three year average for health boards, local authorities and Scotland. Five year average intermediate geographies


Spine Chart

- % =percent
cr2 =crude rate per 1,000 population
cr9 =crude rate per 10,000 population
mean=average
sr4 =age-sex standardised rate per 100,000 population to ESP2013. Please see Appendix I in the technical report.
yrs =years

Spine Chart

Key:

- Statistically significantly 'worse' than National average
- Statistically not significantly different from National average
- Statistically significantly 'better' than National average
- Statistically significant difference compared to National average
- △ No significance can be calculated



See the detailed Definitions and Sources table for indicator information and Technical Report for further guidance on interpreting the spine.

Overview of the ScotPHO profiles for West Lothian

Introduction

This section provides an overview of the population in West Lothian and all the health and wellbeing indicators in the area. The 56 indicators are split into 13 themes (also called domains). For each theme, comparisons are made between what is observed locally and the national picture.

As we include all indicators in the overview summary, some of these differences may be due to random variation. The spine chart (page 4–5) can be used to examine this. The colour of the dot on the spine chart indicates whether the differences are considered to be due to random variation or are statistically significant. Note that an indicator shown as higher does not always mean worse; it could be better. This depends on what the individual indicator measures. Please note that in this overview we use 'similar' wherever a difference is not statistically significant.

Population structure

In 2014, 65% (114,956/177,150) of the population of West Lothian was of working age (16–64 years), similar to the national percentage of 65%. Children and young people (aged 0–15 years) made up 20% (35,064/177,150) of the population, higher than the national 17%. Adults aged over 75 years comprised 6% (10,863/177,150) of the population, lower than the national average of 8%. The population structure of West Lothian has more younger people, fewer older people, and similar people of working age than the national average.

Life expectancy

Life expectancies in 2011, at 77.1 years for males and 80.1 years for females, were similar to the Scottish average of 76.6 years for males and lower than the Scottish female average of 80.8 years.

Mortality

In 2012–2014, the overall mortality rate among young adults (aged 15–44 years) was, at 88, similar to the Scottish rate of 101. Among those aged under 75 years, cancer mortality was 158, so similar to the Scottish rate of 155. For the same age group, for coronary heart disease, the mortality rate was 64, similar to the Scottish rate of 61.

Behaviours

In West Lothian the prevalence of smoking in adults in 2014 was, at 21%, similar to that in Scotland (20%). The rate for smoking-attributable deaths in 2013–2014, at 345, was similar to Scotland (367). The rate for alcohol-related hospital stays in 2014/15 was 642, similar to the rate for Scotland (672). The rate for drug-related hospital stays in 2012/13–2014/15, at 127, was similar to Scotland's 122. In 2010–2014, the rate for alcohol-related deaths, at 20, was similar to the Scottish rate of 23. The percentage of adults walking or cycling to work in 2012/2013, at 10%, was lower than the 16% who did so in Scotland overall.

Ill-health and injury

The rate for cancer registration in 2011–2013 was, at 646, similar to Scotland's overall rate of 634. The rate for patients hospitalised with asthma in 2011–2013, 86, was similar to the Scottish rate of 91. The rate for emergency hospitalisations in 2011–2013, at 8190, was higher than the rate for Scotland (7500). The rate for patients hospitalised for chronic obstructive pulmonary disease (COPD) in 2011–2013, at 710, was higher than the Scottish rate of 660. In 2011–2013, coronary heart disease rate was, at 420, similar to the Scottish level of 440. For road traffic accidents in 2011–2013 the rate was, at 53, similar to the Scottish rate of 63. The rate for adults aged 65 years and over with multiple hospital admissions in 2011–2013, at 5940, was higher than that in Scotland (5160).

Mental health

The percentage of people prescribed medication for anxiety, depression or psychosis in 2014/15 was, at 18%, higher than Scotland overall (17%). The rate for psychiatric hospitalisations in 2011–2013 was 341, which was higher than the Scottish rate of 292. The suicide rate in 2009–2013 was 15, which was similar to the Scottish rate of 15.

Social care and housing

In 2014, 5.1% of adults claimed incapacity benefit, severe disability allowance or employment and support allowance; this was similar to the Scottish figure of 5.1%. The percentage of those aged 65 years and over with high care needs cared for at home, at 38%, was higher than in Scotland overall (35%). The crude rate for children who were looked after by the local authority, at 11/1000, was similar to Scotland's rate of 14/1000. The proportion of households occupied by single adults (35%) was similar to the 38% in Scotland as a whole.

Education

In 2012/13, the mean tariff score – which measures exam success rates – was 206, in comparison with the Scotland mean of 193. In 2010/11, the percentage attendance at primary school in West Lothian, at 95.2%, was higher than Scotland overall (94.8%) and the secondary school attendance of 92% was similar to Scotland's 91%. In 2013, some 11% of working-age adults had low or no educational qualifications, in comparison with 13% in Scotland.

Economy

Economic deprivation indicators suggested that the level of deprivation in West Lothian was similar to the level for Scotland overall. In 2014, the percentage who were income deprived in West Lothian was 12.6% for all ages, lower than the 13.1% for all ages for Scotland. 11.7% were employment deprived, lower than the 12.2% for those employment deprived in Scotland. In 2014, the percentage who claimed out-of-work benefits, at 11.8%, was similar to the 12% across Scotland. In 2014, the percentage of young adults who were outside employment, education or training was 8%, which was higher than the 7% for Scotland. In 2012, the percentage of children living in poverty was 14%, and so was lower than the 15% for Scotland. The percentage of those aged 60 years and over who claimed pension credits in 2014, at 6.9%, was similar to the 6.9% for Scotland.

Crime

In West Lothian the crude population crime rate for 2014 was lower than Scotland at 35/1000 (Scotland: 40/1000) and the crude domestic abuse rate was higher than Scotland at 125/10,000 (Scotland: 112/10,000). The crude rate for recorded drug crimes was lower than Scotland's rate, at 38/10,000 (Scotland: 69/10,000), and the crude rate for referrals to the children's reporter for violence-related offences was compared to Scotland at 3/1000 (Scotland: 2.1/1000). The crude rate for recorded violent crimes was similar to Scotland at 10/10,000 (Scotland: 12/10,000). The rate for prisoner population, at 117, was lower than the Scottish rate of 171.

Environment

In 2013, the percentage of the population who lived within 500m of a derelict site in West Lothian was 25%, which was lower than that in Scotland (30%) and the percentage with access deprivation (i.e. within the 15% of the Scottish population who lived furthest away from local services) was, at 7%, lower than the 15% for Scotland. In 2014, the percentage of adults who rated their area as a very good place to live was, at 61%, similar to Scotland's 56%.

Women's and children's health

In 2011–2013, the crude rate for teenage pregnancy was 41/1000, so similar to Scotland's 41/1000. In 2012/13–2014/15, 2% of births were low weight, similar to Scotland at 2%. In 2013/14 the prevalence of childhood obesity in primary 1 was 8%, so lower than Scotland's 10%. The proportion of mothers smoking in pregnancy, at 22%, was higher than the 19% for Scotland in 2012/13–2014/15. The percentage of exclusive breastfeeding, at 24% in 2012/13–2014/15, was lower than Scotland's 27%. In 2013/14, 64% of children in primary 1 had good dental health, lower than Scotland at 67%. The dental health of children in primary 7, at 43% with no obvious decay experience, was lower than the Scottish average of 48%.

Immunisation and screening

For breast screening, the uptake of 68% in 2010–2012 was lower than the national average of 73%. For bowel screening, the uptake of 53% was lower than the 56% uptake for Scotland. The immunisation uptake for MMR (measles, mumps, rubella) by age 2 years was 96% in 2012–2014, similar to Scotland's 95%. The immunisation uptake for 5 in 1 (diphtheria, pertussis, tetanus, polio, Hib [meningitis]) by age 2 years in 2012–2014 was 99%, higher than Scotland's 98%.

Intermediate zones

Spine charts showing the data that are available for smaller areas within West Lothian (intermediate zones [IZs] – areas with approximately 10,000 individuals) are available via the online tool, but it is not possible to show these in the printed reports owing to the volume of the data (there are 1235 IZs in Scotland). The IZs can be further investigated by selecting Intermediate Zone from the Geography drop-down menu in the 'Profiles update process' section within the tool. This will show a map initially featuring the IZs for Aberdeen City. Use the buttons provided to zoom in (+) or out (–) of the map. You can select any local authority from the drop-down menu to the top-right of the map. To access the spine, trend and rank charts about a particular IZ, choose one from the list in the left-hand drop-down menu. Some indicators, particularly those reliant on survey data or those which have a very small number of outcomes, are not available at this level.

Comparisons, time trends, other indicators and further data

Using the online ScotPHO profile tool (www.scotpho.org.uk/comparative-health/profiles/online-profiles-tool) it is possible to compare the outcomes between any NHS Board, local authority or IZ area. A wider range of outcomes data are also available in the tool, including alcohol and drugs profiles, older adults' profiles, diabetes profiles and mental health profiles.

In any of the ScotPHO profiles the extent to which any differences may be due to random variation or chance is indicated by the colouring of the dots. By clicking on each individual indicator in the online tool it is also possible to compare areas on a rank chart and view time-trend data. Where relevant, data are also presented as standardised rates (to balance and account for age differences between areas) and as crude rates (to reflect the actual number of individuals/events for a particular outcome for service planning). Data across the socioeconomic determinants of health and health topic areas are available on the ScotPHO website: www.scotpho.org.uk

To examine inequalities within local authority areas, use the ScotPHO deprivation profiles from the online tool. These provide data on the inequalities across the local populations broadly in line with the outcomes relevant to Single Outcome Agreements (economic recovery and growth, employment, early years, safer and stronger communities and health inequalities).

A technical report is available on the profiles gateway page (www.scotpho.org.uk/opt/Reports/HWP-2015-technical-report-13112015.pdf), giving more background on the indicators and how they were measured. A user guide for the online profiles tool is also available here.

Contact ScotPHO scotpho@nhs.net for further information.